

# Polar lipid fatty acids, LPS-hydroxy fatty acids, and respiratory quinones of three *Geobacter* strains, and variation with electron acceptor

D. B. Hedrick · A. D. Peacock · D. R. Lovley ·  
T. L. Woodard · K. P. Nevin · P. E. Long · D. C. White

Received: 14 August 2008 / Accepted: 24 September 2008 / Published online: 10 October 2008  
© Society for Industrial Microbiology 2008

**Abstract** The polar lipid fatty acids, lipopolysaccharide hydroxy-fatty acids, and respiratory quinones of *Geobacter metallireducens* str. GS-15, *Geobacter sulfurreducens* str. PCA, and *Geobacter bemidjensis* str. Bem are reported. Also, the lipids of *G. metallireducens* were compared when grown with  $\text{Fe}^{3+}$  or nitrate as electron acceptors and *G. sulfurreducens* with  $\text{Fe}^{3+}$  or fumarate. In all experiments, the most abundant polar lipid fatty acids were 14:0, i15:0, 16:1 $\omega$ 7c, 16:1 $\omega$ 5c, and 16:0; lipopolysaccharide hydroxy-fatty acids were dominated by 3oh16:0, 3oh14:0, 9oh16:0, and 10oh16:0; and menaquinone-8 was the most abundant respiratory quinone. Some variation in lipid profiles with strain were observed, but not with electron acceptor.

**Keywords** *Geobacter* · Polar lipid fatty acids · Lipopolysaccharide fatty acids · Respiratory quinones · Lipid analysis

---

D. C. White: Deceased.

---

D. B. Hedrick (✉)  
Microbial Insights, Inc., 2340 Stock Creek Blvd.,  
Rockford, TN, USA  
e-mail: davidhedrick@earthlink.net

A. D. Peacock  
Haley & Aldrich, 465 Medford St., Ste. 2200,  
Boston, MA, USA

D. R. Lovley · T. L. Woodard · K. P. Nevin  
Department of Microbiology, University of Massachusetts,  
159 N. Pleasant St., Amherst, MA, USA

P. E. Long  
Pacific Northwest National Laboratory, Richland, WA, USA

D. C. White  
Center for Biomarker Analysis, University of Tennessee,  
10515 Research Dr., Ste. 300, Knoxville, TN, USA

## Abbreviations

LPS ohFA	Lipopolysaccharide hydroxy-fatty acids
PLFA	Polar lipid fatty acids
MK	Menaquinone

## Introduction

*Geobacter* is a versatile Deltaproteobacterium that can utilize  $\text{Fe}^{3+}$ ,  $\text{Mn}^{4+}$ ,  $\text{U}^{6+}$  [18], or oxygen [14] as electron acceptors which has important roles in geochemical cycling [18], bioremediation [25, 32], and electrical generation [30]. In support of ecological studies using  $^{13}\text{C}$ -labeled substrate incorporation into lipid biomarkers of *Geobacter*, and general biochemical studies, the polar lipid fatty acids (PLFA), lipopolysaccharide hydroxy-fatty acids (LPS ohFA), and respiratory quinones of three strains of *Geobacter* were determined, and on two of the strains with different electron acceptors.

## Materials and methods

### Strains and culturing

*Geobacter metallireducens* str. GS-15 [18], *Geobacter sulfurreducens* str. PCA [4], and *Geobacter bemidjensis* str. Bem [23] were cultured in freshwater medium [19] with 55 mM  $\text{Fe}^{3+}$  citrate as electron acceptor and 10 mM acetate as electron donor. *G. metallireducens* was also cultured in the same medium with 5 mM nitrate replacing  $\text{Fe}^{3+}$  citrate as electron acceptor, and *G. sulfurreducens* was cultured in NB medium with 40 mM fumarate as electron acceptor and 15 mM acetate as electron donor [17]. All culturing was in batch at 30°C under standard anaerobic conditions [1, 26].

Cells were harvested by centrifugation at 3,200g for 20 min and frozen at  $-20^{\circ}\text{F}$  until analysis.

#### Polar lipid fatty acid analysis

Dried cell material was extracted using the method of Bligh and Dyer [3] as modified [29]. Complete extraction was ensured by sonication for 2 min and extraction for 4 h before the separation of the aqueous and organic phases. The total lipid extract was then fractionated into neutral lipid, glycolipid, and phospholipid by elution with chloroform, acetone, and methanol, respectively, using 500 mg silicic acid columns [10]. Neutral lipids were reserved for quinone analysis. The polar lipids were transesterified to the fatty acid methyl esters using a mild alkaline methanolysis [10].

An Agilent 6890 series gas chromatograph with a 50 m nonpolar column (0.2 mm I.D., 0.11  $\mu\text{m}$  film thickness) was used for gas chromatographic analysis. The temperature program was initial temperature =  $100^{\circ}\text{C}$ ,  $10^{\circ}\text{C}/\text{min}$  to  $150^{\circ}\text{C}$ ,  $150^{\circ}\text{C}$  for 1 min,  $3^{\circ}\text{C}/\text{min}$  to  $282^{\circ}\text{C}$ , and hold at  $282^{\circ}\text{C}$  for 5 min. The injector and detector temperatures were 270 and  $290^{\circ}\text{C}$ , respectively. Peak quantitation and mass identification was performed using an Agilent 5973 mass-selective detector. Monounsaturates were identified by derivatization with dimethyldisulfide and gas chromatography/mass spectroscopy as described [24].

#### Lipopolysaccharide hydroxy-fatty acid analysis

Dried cell material was treated with a strong acid methanolysis (chloroform/methanol/hydrochloric acid 10:1:1 at  $100^{\circ}\text{C}$  for 1 h) to liberate the fatty acids as their methyl esters [12]. The hydroxy moieties were converted to trimethylsilyl ethers and analyzed by the same instrumentation and methods as the polar lipid fatty acids. Only the hydroxy-fatty acids were reported.

#### Respiratory quinone analysis

The respiratory quinones in the neutral lipid fraction of the total lipid extract were determined by atmospheric pressure chemical ionization tandem mass spectrometry as described [9].

## Results

#### Polar Lipid Fatty Acids

The PLFA profiles (Table 1) of the *Geobacter* strains analyzed were very similar to each other, with 14:0, i15:0, 16:1 $\omega$ 7c, 16:1 $\omega$ 5c, and 16:0 more than 1% of the total for

all strains under all conditions. 18:1 $\omega$ 7c was also more than 1% in some samples. The biggest differences between strains were in i15:0 and 18:1 $\omega$ 7c (*G. metallireducens* > *G. sulfurreducens* > *G. bemidjensis*). The effect of electron acceptor on the PLFA profiles were less than the differences between the strains. *G. metallireducens* with nitrate as the electron acceptor and *G. sulfurreducens* with fumarate had slightly higher proportions of 16:1 $\omega$ 7c and less 16:0 compared to with ferric iron as the electron acceptor.

#### Lipopolysaccharide hydroxy-fatty acids

The LPS oh-FA (Table 1) were more variable between strains and electron acceptors than the PLFA. *G. metallireducens* was distinguished by having over four times more 3oh16:0 than 3oh14:0, and significant amounts of 9oh16:0 and 10oh16:0 (10–15%), while *G. sulfurreducens* and *G. bemidjensis* had more similar proportions of 3oh16:0 and 3oh14:0 and much less 9oh16:0 and 10oh16:0 (5–0.4%). As was found for the PLFA, the variation between strains was greater than the variation with electron acceptor.

#### Respiratory quinones

The respiratory quinone profiles (Table 1) were dominated by menaquinone-8 (MK-8), which was more than 85% of the total quinones for all strains under all conditions. Ubiquinones were not detected above background. Overall, the quinone profiles were very similar to each other between strains and treatments.

## Discussion

The PLFA and LPS ohFA composition of *G. metallireducens* under iron- and nitrate-reducing conditions has been determined [18]. While the overall lipid profiles were very similar, the shifts in the profiles with change in electron acceptor were different. In this work (Table 1) all of the major PLFA (>1%) decreased except for 16:1 $\omega$ 7c when the electron acceptor was changed from ferric iron to nitrate. However, in the experiment performed over 10 years earlier, 14:0 and i15:0 increased. Similarly for the LPS oh-FA, in this work 3oh14:0, 9oh16:0, and 10oh16:0 increased, while in [18] 3oh14:0 and 3oh15:0 increased. *G. metallireducens* grown with  $\text{Fe}^{3+}$  citrate had similar overall profiles, except that 18:1 $\omega$ 7c was 29% [36], up from 1.1 to 3.7% in this study and [18]. This suggests that the PLFA and LPS oh-FA of *G. metallireducens* are less affected by the change in electron acceptor, more by details of culturing conditions. Growth phase, ionic strength, temperature, toxic compounds, and carbon source have all been shown to shift bacterial fatty acid profiles.

**Table 1** Polar lipid fatty acids, lipopolysaccharide hydroxy-fatty acids, and respiratory quinones of three strains of *Geobacter* with different electron acceptors, as mole percents, average and standard deviation ( $n = 3$ )

Strain	<i>Geobacter metallireducens</i> , Avg (SD)		<i>Geobacter sulfurreducens</i> , Avg (SD)		<i>G. bemidjiensis</i> , Avg (SD)
	Fe <sup>3+</sup>	Nitrate	Fe <sup>3+</sup>	Fumarate	
Polar lipid fatty acids					
14:1 $\omega$ 7c D	0.04 (0.03)	0.03 (0.05)	0.20 (0.03)	0.28 (0.01)	0.38 (0.08)
14:1 $\omega$ 7t	0.05 (0.04)	0.03 (0.05)	0.07 (0.02)	0.08 (0.00)	0.09 (0.02)
14:1 $\omega$ 5c					0.16 (0.05)
<b>14:0</b>	<b>6.39</b> (0.86)	<b>6.23</b> (5.10)	<b>9.19</b> (0.54)	<b>8.49</b> (0.14)	<b>7.67</b> (0.34)
i15:1 $\omega$ 7c	0.05 (0.04)	0.02 (0.03)	0.10 (0.02)	0.11 (0.01)	
<b>i15:0</b>	<b>9.88</b> (0.27)	<b>9.66</b> (2.26)	<b>6.31</b> (0.40)	<b>5.32</b> (0.11)	<b>1.07</b> (0.04)
a15:0	0.27 (0.05)	0.05 (0.09)	0.54 (0.06)	0.27 (0.01)	0.14 (0.04)
15:1				0.01 (0.01)	0.13 (0.04)
15:0			0.61 (0.74)	0.20 (0.02)	0.14 (0.03)
<b>16:1<math>\omega</math>7c D</b>	<b>48.44</b> (1.08)	<b>57.40</b> (3.36)	<b>35.87</b> (0.55)	<b>38.30</b> (0.20)	<b>46.02</b> (0.79)
16:1 $\omega$ 7t D	0.74 (0.12)	0.91 (0.23)	0.41 (0.09)	0.39 (0.05)	0.70 (0.02)
<b>16:1<math>\omega</math>5c D</b>	<b>1.65</b> (0.10)	<b>1.49</b> (0.36)	<b>1.11</b> (0.07)	<b>1.41</b> (0.03)	<b>2.22</b> (0.01)
<b>16:0</b>	<b>29.27</b> (0.37)	<b>22.66</b> (3.76)	<b>43.52</b> (0.63)	<b>42.69</b> (0.19)	<b>40.56</b> (0.78)
i17:1 $\omega$ 7c D	0.13 (0.02)	0.09 (0.08)	0.14 (0.04)	0.11 (0.02)	
10Me16:0	0.46 (0.06)	0.19 (0.07)	0.18 (0.06)	0.12 (0.06)	0.06 (0.06)
i17:0	0.25 (0.04)	0.08 (0.07)	0.08 (0.02)	0.07 (0.01)	
<b>18:1<math>\omega</math>7c D</b>	<b>2.00</b> (0.19)	0.94 (0.34)	<b>1.13</b> (0.10)	<b>1.24</b> (0.03)	0.48 (0.07)
18:1 $\omega$ 5c	0.09 (0.03)		0.05 (0.00)	0.08 (0.01)	
18:0	0.28 (0.06)	0.19 (0.06)	0.30 (0.03)	0.52 (0.01)	0.14 (0.01)
br19:1			0.06 (0.00)	0.09 (0.01)	
Lipopolysaccharide hydroxy-fatty acids					
3oh12:0			0.7 (1.1)	1.3 (1.3)	0.2 (0.1)
3oh13:0					0.1 (0.1)
<b>3oh14:0</b>	<b>11.1</b> (1.9)	<b>13.4</b> (2.4)	<b>30.0</b> (4.1)	<b>46.9</b> (8.5)	<b>47.8</b> (1.9)
3oh-i15:0			4.4 (0.4)	7.6 (1.4)	7.1 (0.1)
3oh-a15:0				1.3 (2.2)	0.4 (0.1)
3oh15:0	6.7 (5.8)		1.3 (2.3)		0.6 (0.1)
<b>3oh16:0</b>	<b>62.3</b> (6.1)	<b>55.9</b> (10.7)	<b>53.5</b> (4.1)	<b>38.2</b> (5.9)	<b>42.8</b> (1.8)
<b>9oh16:0</b>	<b>9.0</b> (1.6)	<b>14.1</b> (5.7)	4.5 (0.2)	2.2 (1.9)	0.4 (0.3)
<b>10oh16:0</b>	<b>11.0</b> (1.7)	<b>16.6</b> (6.4)	5.5 (0.3)	2.5 (2.3)	0.6 (0.0)
Respiratory quinones					
MK-4	3.39 (0.15)	3.28 (0.21)	2.77 (0.13)	3.39 (0.55)	3.07 (0.07)
MK-5	1.52 (0.11)	1.50 (0.06)	1.36 (0.02)	1.41 (0.09)	1.48 (0.06)
MK-6	4.73 (0.06)	4.68 (0.08)	4.35 (0.09)	5.30 (1.00)	4.61 (0.04)
MK-7	0.68 (0.04)	0.93 (0.06)	0.66 (0.02)	0.77 (0.17)	1.21 (0.01)
<b>MK-8</b>	<b>85.18</b> (0.36)	<b>86.86</b> (0.16)	<b>87.17</b> (0.16)	<b>86.69</b> (1.67)	<b>87.21</b> (0.25)
MK-9	4.45 (0.33)	2.73 (0.18)	3.68 (0.14)	2.44 (0.14)	2.41 (0.17)

Major components (fatty acids > 1%, LPS oh-FA and MK > 10%) are in bold. Fatty acids marked with a “D” were validated by DMDS derivatization. Fatty acids not greater than 0.1% in any analysis were not reported, including i14:0, 3oh14:0, i16:0, a16:0, 11Me16:0, Cy17:0, 17:0, br18:1, 3oh16:0, and 18:1 $\omega$ 7t

The PLFA profiles distinguished *G. metallireducens*, *sulfurreducens*, and *bemidjiensis* in pure culture under controlled conditions, but could not be used to determine

which organism is present in the complex microbial community of an environmental sample. It will also be difficult to distinguish some Deltaproteobacterial sulfate-reducers

from *Geobacter* in environmental samples by PLFA analysis, unless *Geobacter* is a high proportion of the community. *Desulfuromonas acetoxydans* [6], a co-member of the Desulfuromonadales with *Geobacter*, has a very similar profile to the *Geobacters* discussed above. Desulfovibrionales have more iso- and anteiso-branched 15- and 17-carbon fatty acids and i17:1 $\omega$ 7c [7, 8, 15, 34]. Desulfobacteraceae have more 10Me16:0 and Cy17:0 [6, 15, 34]. However, in the case of subsurface biostimulation of U(VI) immobilization by *Geobacter*, the high cell numbers of *Geobacter* in treated sediments allows monitoring its population by PLFA analysis [22].

The LPS ohFA could be used to detect the growth of *Geobacter* in the subsurface during bioremediation. The strains analyzed had unusually high levels of 3oh16:0 (Table 1), and the unusual fatty acids 9oh16:0 and 10oh16:0. The literature on LPS FA is incomplete, especially for environmental isolates, but a study of 6 strains of *Desulfovibrio* found 3oh-i15:0 or 3oh-i17:0 as the most abundant LPSFA [7]. 3oh16:0 was only 5.8–16.6% in these strains. A saturated subsurface sediment sample had 16.9% 3oh16:0 in the LPSFA, and no 3oh16:0 in any of the 4 Gammaproteobacterial strains tested [27]. Since LPSFA are not degraded as quickly as PLFA upon cell death, sediment samples can have much more LPSFA than can be accounted for by the viable microbial biomass, representing “fossil” bacteria (Hedrick and White, unpublished). Therefore, groundwater or Bio-Trap<sup>®</sup> samples [28] would be more appropriate for this analysis.

Menaquinone-8 accounted for over 85% of the menaquinones and no ubiquinones were detected for all three strains and two redox acceptor experiments. MK-8 has been detected before in *G. sulfurreducens* [21]. The high proportion of this single respiratory quinone in *Geobacter* makes it a possible candidate as a biomarker. Anaerobic Gammaproteobacteria and Bacteroidetes often have MK-8, [5], but the sulfate-reducers usually do not. MK-6 has been reported from *Desulfovibrio desulfuricans*, *Desulfovibrio gigas*, and *Desulfovibrio vulgaris* [11, 20, 35]; MK-7 in *Desulfovermiculus halophilus* [2]; MK-5(H2) in *Desulfobulbus rhabdiformis* [13]; MK-7(H2) in *Desulfovibrio magneticus* [31]; no menaquinones were detected in *Desulfomonile tiedjei* DCB-1 [16]; and only *Desulfopila aestuarii* had MK-8 [33].

**Acknowledgments** This research was supported by the Office of Science of the US Department of Energy, Biological and Environmental Research, grant DE-FG02-04ER63719.

## References

- Balch WE, Fox GE, Magrum LJ, Woese CR, Wolfe RS (1979) Methanogens: reevaluation of a unique biological group. *Microbiol Rev* 43:260–296
- Belyakova E, Rozanova E, Borzenkov I, Tourova T, Pusheva M, Lysenko A et al (2006) The new facultatively chemolithoautotrophic, moderately halophilic, sulfate-reducing bacterium *Desulfovermiculus halophilus* gen. nov., sp. nov., isolated from an oil field. *Microbiology* 75:161–171. doi:10.1134/S0026261706020093
- Bligh EG, Dyer WJ (1959) A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 37:911–917
- Caccavo F, Lonergan DJ, Lovley DR, Davis M, Stolz JF, McInerney MJ (1994) *Geobacter sulfurreducens* sp. nov., a hydrogen- and acetate-oxidizing dissimilatory metal-reducing microorganism. *Appl Environ Microbiol* 60:3752–3759
- Collins MD, Jones D (1981) Distribution of isoprenoid quinone structural types in bacteria and their taxonomic implications. *Microbiol Rev* 45:316–354
- Dowling NJE, Widdel F, White DC (1986) Phospholipid ester-linked fatty acid biomarkers of acetate-oxidizing sulfate reducers and other sulfide forming bacteria. *J Gen Microbiol* 132:1815–1825
- Edlund A, Nichols PD, Roffky R, White DC (1985) Extractable and lipopolysaccharide fatty acid and hydroxy acid profiles from *Desulfovibrio* species. *J Lipid Res* 26:982–988
- Feio MJ, Beech IB, Carepo M, Lopes JM, Cheung CWS, Franco R et al (1998) Isolation and characterization of a novel sulphate-reducing bacterium of the *Desulfovibrio* genus. *Anaerobe* 4:117–130. doi:10.1006/anae.1997.0142
- Geyer R, Peacock AD, White DC, Lytle C, Van Berkel GJ (2004) Atmospheric pressure chemical ionization and atmospheric pressure photoionization for simultaneous mass spectrometric analysis of microbial respiratory ubiquinones and menaquinones. *J Mass Spectrom* 39:922–929. doi:10.1002/jms.670
- Guckert JB, Antworth CP, Nichols PD, White DC (1985) Phospholipid, ester-linked fatty acid profiles as reproducible assays for changes in prokaryotic community structure of estuarine sediments. *FEMS Microbiol Ecol* 31:147–158
- Hatchikian EC (1974) On the role of menaquinone-6 in the electron transport of hydrogen: fumarate reductase system in the strict anaerobe *Desulfovibrio gigas*. *J Gen Microbiol* 81:261–266
- Hedrick DB, Guckert JB, White DC (1991) Archaeobacterial ether lipid diversity analyzed by supercritical fluid chromatography: integration with a bacterial lipid protocol. *J Lipid Res* 32:659–666
- Lien T, Madsen M, Steen IH, Gjerdevik K (1998) *Desulfobulbus rhabdiformis* sp. nov., a sulfate reducer from a water–oil separation system. *Int J Syst Bacteriol* 48:469–474
- Lin WC, Coppi MV, Lovley DR (2004) *Geobacter sulfurreducens* can grow with oxygen as a terminal electron acceptor. *Appl Environ Microbiol* 70:2525–2528. doi:10.1128/AEM.70.4.2525-2528.2004
- Londry KL, Jahnke LL, Des Marais DJ (2004) Stable carbon isotope ratios of lipid biomarkers of sulfate-reducing bacteria. *Appl Environ Microbiol* 70:745–751. doi:10.1128/AEM.70.2.745-751.2004
- Louie TM, Mohn WW (1999) Evidence for a chemiosmotic model of dehalorespiration in *Desulfomonile tiedjei* DCB-1. *J Bacteriol* 181:40–46
- Lovley DR, Fraga JL, Coates JD, Blunt-Harris EL (1999) Humics as an electron donor for anaerobic respiration. *Environ Microbiol* 1:89–98. doi:10.1046/j.1462-2920.1999.00009.x
- Lovley DR, Giovannoni SJ, White DC, Champine JE, Phillips EJP, Gorby YA et al (1993) *Geobacter metallireducens* gen. nov. sp. nov., a microorganism capable of coupling the complete oxidation of organic compounds to the reduction of iron and other metals. *Arch Microbiol* 159:336–344
- Lovley DR, Phillips EJP (1988) Novel mode of microbial energy metabolism: organic carbon oxidation coupled to dissimilatory reduction of iron or manganese. *Appl Environ Microbiol* 54:1472–1480

20. Maroc J, Azerad R, Kamen D, Gall JL (1970) Menaquinone (MK-6) in the sulfate-reducing obligate anaerobes, *Desulfovibrio*. Biochim Biophys Acta 197:87–89. doi:[10.1016/0005-2728\(70\)90012-5](https://doi.org/10.1016/0005-2728(70)90012-5)
21. Mikoulinskaia O, Akimenko V, Galouchko A, Thauer RK, Hedderich R (1999) Cytochrome c-dependent methacrylate reductase from *Geobacter sulfurreducens* AM-1. Eur J Biochem 263:346–352. doi:[10.1046/j.1432-1327.1999.00489.x](https://doi.org/10.1046/j.1432-1327.1999.00489.x)
22. Mohanty SR, Kollah B, Hedrick DB, Peacock AD, Kukkadapu RK, Roden EE (2008) Biogeochemical processes in ethanol stimulated uranium-contaminated subsurface sediments. Environ Sci Technol 42:4384–4390. doi:[10.1021/es703082v](https://doi.org/10.1021/es703082v)
23. Nevin KP, Holmes DE, Woodard TL, Hinlein ES, Ostendorf DW, Lovley DR (2005) *Geobacter bemidjensis* sp. nov. and *Geobacter psychrophilus* sp. nov., two novel Fe(III)-reducing subsurface isolates. Int J Syst Evol Microbiol 55:1667–1674. doi:[10.1099/ijs.0.63417-0](https://doi.org/10.1099/ijs.0.63417-0)
24. Nichols PD, Guckert JB, White DC (1986) Determination of monounsaturated fatty acid double-bond position and geometry for microbial monocultures and complex consortia by capillary GC-MS of their dimethyl disulphide adducts. J Microbiol Methods 5:49–55. doi:[10.1016/0167-7012\(86\)90023-0](https://doi.org/10.1016/0167-7012(86)90023-0)
25. North NN, Dollhopf SL, Petrie L, Istok JD, Balkwill DL, Kostka JE (2004) Change in bacterial community structure during in situ biostimulation of subsurface sediment cocontaminated with uranium and nitrate. Appl Environ Microbiol 70:4911–4920. doi:[10.1128/AEM.70.8.4911-4920.2004](https://doi.org/10.1128/AEM.70.8.4911-4920.2004)
26. Nottingham PM, Hungate RE (1969) Methanogenic fermentation of benzoate. J Bacteriol 98:1170–1172
27. Parker JH, Smith GA, Fredrickson HL, Vestal JR, White DC (1982) Sensitive assay, based on hydroxy fatty acids from lipopolysaccharide lipid A, for gram-negative bacteria in sediments. Appl Environ Microbiol 44:1170–1177
28. Peacock AD, Chang Y-J, Istok JD, Krumholz L, Geyer R, Kinsall B et al (2004) Utilization of microbial biofilms as monitors of bioremediation. Microb Ecol 47:284–292. doi:[10.1007/s00248-003-1024-9](https://doi.org/10.1007/s00248-003-1024-9)
29. Peacock AD, Mullen MD, Ringelberg DB, Tyler DD, Hedrick DB, Gale PM et al (2000) Soil microbial community responses to dairy manure or ammonium nitrate applications. Soil Biol Biochem 33:1011–1019. doi:[10.1016/S0038-0717\(01\)00004-9](https://doi.org/10.1016/S0038-0717(01)00004-9)
30. Reguera G, Nevin KP, Nicoll JS, Covalla SF, Woodard TL, Lovley DR (2006) Biofilm and nanowire production lead to increased current in microbial fuel cells. Appl Environ Microbiol 72:7345–7348. doi:[10.1128/AEM.01444-06](https://doi.org/10.1128/AEM.01444-06)
31. Sakaguchi T, Arakaki A, Matsunaga T (2002) *Desulfovibrio magneticus* sp. nov., a novel sulfate-reducing bacterium that produces intracellular single-domain-sized magnetite particles. Int J Syst Evol Microbiol 52:215–221
32. Sung Y, Fletcher KE, Ritalahti KM, Apkarian RP, Ramos-Hernández N, Sanford RA et al (2006) *Geobacter lovleyi* sp. nov. strain SZ, a novel metal-reducing and tetrachloroethene-dechlorinating bacterium. Appl Environ Microbiol 72:2775–2782. doi:[10.1128/AEM.72.4.2775-2782.2006](https://doi.org/10.1128/AEM.72.4.2775-2782.2006)
33. Suzuki D, Ueki A, Amaishi A, Ueki K (2007) *Desulfopila aestuarii* gen. nov., sp. nov., a Gram-negative, rod-like, sulfate-reducing bacterium isolated from an estuarine sediment in Japan. Int J Syst Evol Microbiol 57:520–526. doi:[10.1099/ijs.0.64600-0](https://doi.org/10.1099/ijs.0.64600-0)
34. Taylor J, Parkes RJ (1983) The cellular fatty acids of the sulphate-reducing bacteria, *Desulfobacter* sp., *Desulfobulbus* sp. and *Desulfovibrio desulfuricans*. J Gen Microbiol 129:3303–3309
35. Weber MM, Matachiner JT, Peck HD (1970) Menaquinone-6 in the strict anaerobes *Desulfovibrio vulgaris* and *Desulfovibrio gigas*. Biochem Biophys Res Commun 38:197–204. doi:[10.1016/0006-291X\(70\)90696-0](https://doi.org/10.1016/0006-291X(70)90696-0)
36. Zhang CL, Li Y, Ye Q, Fong J, Peacock AD, Blunt E et al (2003) Carbon isotope signatures of fatty acids in *Geobacter metallireducens* and *Shewanella algae*. Chem Geol 195:17–28. doi:[10.1016/S0009-2541\(02\)00386-8](https://doi.org/10.1016/S0009-2541(02)00386-8)